

Boundedness commutator between Sobolev spaces

By Sadek Gala
 Université d'Evry Val d'Essonne
 Département de mathématiques
 Bd F. Mitterrand. 91025 Evry Cedex. France
 Sadek.Gala@maths.univ-evry.fr

1st February 2008

Abstract

In this paper, we will study the boundedness properties of commutator

$$C_f = [f, \Delta]$$

acting from $\dot{H}^1(\mathbb{R}^d)$ to $\dot{H}^{-1}(\mathbb{R}^d)$.

Throughout this paper, d will denote a fixed positive integer greater than 2 and Ω will denote an open, non-empty subset of $\mathbb{R}^d$. We wish to study linear elliptic differential operator Δ in divergence form defined on C^2 functions by

$$\Delta f(x) = \frac{1}{2} \sum_{i,j=1}^d \partial_i (\partial_j f)(x)$$

Recall the divergence theorem. Suppose Ω is a nice region, F is a smooth vector field, $\nu(x)$ is the outward normal vector at $x \in \partial\Omega$, and σ is surface measure on $\partial\Omega$. The divergence theorem then says that

$$\int_{\partial\Omega} F(y) \cdot \nu(y) \sigma(dy) = \int_{\Omega} \operatorname{div} F(x) dx.$$

A twice continuously differentiable, complex-valued function v defined on Ω is harmonic on Ω if

$$\Delta v = 0.$$

We begin with the following simple observation :

Proposition 0.1 *Let g be a C^∞ function with compact support and f a bounded C^∞ function. Then*

$$\int_{\mathbb{R}^d} g(x) \Delta f(x) dx = -\frac{1}{2} \int_{\mathbb{R}^d} \left(\sum_{i,j=1}^d \partial_i g(x) \partial_j f(x) \right) dx.$$

The integrand on the right could be written $\nabla g \cdot \nabla f$.

Proof. We apply the divergence theorem. Let B be a ball large enough to contain the support of g and let $F(x)$ be the vector field whose i -th component is

$$\frac{1}{2} g(x) \sum_{j=1}^d \partial_j f(x).$$

Since g is 0 on ∂B , then $F \cdot \nu = 0$ on ∂B , and also,

$$\operatorname{div} F(x) = \frac{1}{2} \sum_{i=1}^d \partial_i \left(g(x) \sum_{j=1}^d \partial_j f(x) \right) = \frac{1}{2} \sum_{i,j=1}^d \partial_i g(x) \partial_j f(x) + g(x) \Delta f(x).$$

Proposition 0.2 *Suppose v is positive and harmonic in $Q(4)$. There exists c_1 independent of v such that if $w = \log v$, then*

$$\int_Q |\nabla w(x)|^2 dx \leq c_1 h^{d-2}$$

for all cubes Q of side length h contained in $Q(2)$.

Proof. Let Q^* be the cube with the same center as Q but side length twice as long. Note $Q^* \subset Q(4)$. Let φ be C^∞ with values in $[0, 1]$, equal to 1 on Q , supported in Q^* , and such that $\|\nabla \varphi\|_\infty \leq c_2 h^{-1}$. Since

$$\nabla w = \frac{\nabla v}{v}$$

and v is harmonic in $Q(4)$,

$$\begin{aligned} 0 &= 2 \int \frac{\varphi^2}{v} \Delta v = - \int \nabla \left(\frac{\varphi^2}{v} \right) \cdot \nabla v = - \int 2 \frac{\varphi \nabla \varphi}{v} \cdot \nabla v + \int \frac{\varphi^2}{v^2} \nabla v \cdot \nabla v \\ &= -2 \int \varphi \nabla \varphi \cdot \nabla w + \int \varphi^2 \nabla w \cdot \nabla w. \end{aligned}$$

So by the Cauchy-Schwarz inequality and proposition 0.1,

$$\begin{aligned} \int_{Q^*} \varphi^2 |\nabla w|^2 dx &\leq c_3 \int_{Q^*} \varphi^2 \nabla w \cdot \nabla w = c_4 \int_{Q^*} \nabla \varphi \cdot \varphi \nabla w \\ &\leq c_5 \left(\int_{Q^*} |\nabla \varphi|^2 dx \right)^{\frac{1}{2}} \left(\int_{Q^*} \varphi^2 |\nabla w|^2 dx \right)^{\frac{1}{2}}. \end{aligned}$$

Dividing by the second factor on the right, squaring and using the bound on $|\nabla \varphi|$,

$$\int_Q |\nabla w(x)|^2 dx \leq \left(\int_{Q^*} \varphi^2 |\nabla w|^2 dx \right) \leq c_5^2 |Q^*| c_2^2 h^{-2},$$

which implies our result.

We need to distinguish the class of vector fields $\vec{F}$ such that the commutator inequality

$$\left| \int_{\mathbb{R}^d} \vec{F} \cdot (\bar{u} \nabla v - v \nabla \bar{u}) dx \right| \leq C \|u\|_{\dot{H}^1} \|v\|_{\dot{H}^1}$$

for all $u, v \in \mathcal{D}(\mathbb{R}^d)$. In the important case where $\vec{F} = \nabla f$, the preceding inequality is equivalent to the boundedness of the commutator

$$C_f = [f, \Delta]$$

acting from $\dot{H}^1(\mathbb{R}^d)$ to $\dot{H}^{-1}(\mathbb{R}^d)$.

Another useful point in the sequel is the symmetry of the Laplace operator,

$$\langle \Delta \varphi, \psi \rangle = - \langle D \varphi, D \psi \rangle = \langle \varphi, \Delta \psi \rangle$$

for all $\varphi, \psi \in \mathcal{D}(\mathbb{R}^d)$.

Let us present a proof of the

Lemma 0.3 For any twice continuously differentiable functions $u : A \rightarrow \mathbb{R}$ and $v : A \rightarrow \mathbb{R}$, where $A \subset \mathbb{R}^d$, we have

$$\operatorname{div} (u \nabla v - v \nabla u) = u \Delta v - v \Delta u.$$

Proof. Note that

$$\begin{aligned} \operatorname{div} (u \nabla v - v \nabla u) &= \operatorname{div} \left(u \left(\frac{\partial v}{\partial x_1}, \dots, \frac{\partial v}{\partial x_d} \right) - v \left(\frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_d} \right) \right) \\ &= \operatorname{div} \left(u \frac{\partial v}{\partial x_1} - v \frac{\partial u}{\partial x_1}, \dots, u \frac{\partial v}{\partial x_d} - v \frac{\partial u}{\partial x_d} \right) \\ &= \frac{\partial}{\partial x_1} \left(u \frac{\partial v}{\partial x_1} - v \frac{\partial u}{\partial x_1} \right) + \dots + \frac{\partial}{\partial x_d} \left(u \frac{\partial v}{\partial x_d} - v \frac{\partial u}{\partial x_d} \right) \\ &= \left(u \frac{\partial^2 v}{\partial x_1^2} + \frac{\partial u}{\partial x_1} \frac{\partial v}{\partial x_1} - \frac{\partial v}{\partial x_1} \frac{\partial u}{\partial x_1} - v \frac{\partial^2 u}{\partial x_1^2} \right) + \dots + \left(u \frac{\partial^2 v}{\partial x_d^2} + \frac{\partial u}{\partial x_d} \frac{\partial v}{\partial x_d} - \frac{\partial v}{\partial x_d} \frac{\partial u}{\partial x_d} - v \frac{\partial^2 u}{\partial x_d^2} \right) \\ &= u \left(\frac{\partial^2 v}{\partial x_1^2} + \dots + \frac{\partial^2 v}{\partial x_d^2} \right) - v \left(\frac{\partial^2 u}{\partial x_1^2} + \dots + \frac{\partial^2 u}{\partial x_d^2} \right) = u \Delta v - v \Delta u. \end{aligned}$$

Lemma 0.4 Let $f \in \mathcal{D}'(\mathbb{R}^d)$. Then for all $u, v \in \mathcal{D}(\mathbb{R}^d)$, we have

$$|\langle C_f u, v \rangle| = \left| \int_{\mathbb{R}^d} \vec{F} \cdot (\bar{u} \nabla v - v \nabla \bar{u}) dx \right|.$$

Proof. Let $u, v \in \mathcal{D}(\mathbb{R}^d)$. We observe that

$$\begin{aligned} \langle C_f u, v \rangle &= \langle [f, \Delta] u, v \rangle = \langle f \Delta u - \Delta (fu), v \rangle \\ &= \langle f \Delta u, v \rangle - \langle \Delta (fu), v \rangle = \langle f, v \Delta \bar{u} \rangle - \langle fu, \Delta v \rangle \\ &= \langle f, v \Delta \bar{u} \rangle - \langle f, \bar{u} \Delta v \rangle = \langle f, v \Delta \bar{u} - \bar{u} \Delta v \rangle \\ &= \langle f, \operatorname{div} (v \nabla \bar{u} - \bar{u} \nabla v) \rangle \\ &= - \langle \nabla f, v \nabla \bar{u} - \bar{u} \nabla v \rangle = - \left\langle \vec{F}, v \nabla \bar{u} - \bar{u} \nabla v \right\rangle \end{aligned}$$

Therefore

$$|\langle C_f u, v \rangle| = \left| \int_{\mathbb{R}^d} \vec{F} \cdot (\bar{u} \nabla v - v \nabla \bar{u}) dx \right|.$$

We will let P be the Newton potential of μ defined by

$$P\mu(x) = P(x) = C(d) \int_{\mathbb{R}^d} \frac{d\mu(y)}{|x-y|^{d-2}}, \quad x \in \mathbb{R}^d, \quad d \geq 3.$$

Then, we have the following lemma :

Lemma 0.5 *Let μ be a positive Borel measure on $\mathbb{R}^d$ such that $P(x) = I_2\mu(x) \neq \infty$. Then the following inequality hold :*

$$\int_{\mathbb{R}^d} u^2(x) \frac{|\nabla P(x)|^2}{P^2(x)} dx \leq 4 \|\nabla u\|_{L^2}^2, \quad u \in \mathcal{D}(\mathbb{R}^d). \quad (1)$$

Proof. Suppose $u \in \mathcal{D}(\mathbb{R}^d)$. Then $K = \text{supp } u$ is a compact set, and obviously $\inf_{x \in K} P(x) > 0$. Without loss of generality, we assume that $\nabla P \in L_{loc}^2(\mathbb{R}^d)$, and hence the left-hand side of (1) is finite.

Using integration by parts, together with the properties $-\Delta P = \mu$ (understood in the distributional sense) and applying the Schwarz inequality, we get :

$$\begin{aligned} 2 \int_{\mathbb{R}^d} \nabla u(x) \cdot \nabla P(x) \frac{u(x)}{P(x)} dx &= \int_{\mathbb{R}^d} u^2(x) \frac{|\nabla P(x)|^2}{P^2(x)} dx + \int_{\mathbb{R}^d} u^2(x) \frac{d\mu(x)}{P(x)} dx \\ &\leq 2 \left(\int_{\mathbb{R}^d} u^2(x) \frac{|\nabla P(x)|^2}{P^2(x)} dx \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^d} |\nabla u(x)|^2 dx \right)^{\frac{1}{2}}, \end{aligned}$$

for all $u \in \mathcal{D}(\mathbb{R}^d)$. Using this inequality, we have

$$\int_{\mathbb{R}^d} u^2(x) \frac{|\nabla P(x)|^2}{P^2(x)} dx \leq 2 \left(\int_{\mathbb{R}^d} u^2(x) \frac{|\nabla P(x)|^2}{P^2(x)} dx \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^d} |\nabla u(x)|^2 dx \right)^{\frac{1}{2}}.$$

Consequently, we obtain

$$\int_{\mathbb{R}^d} u^2(x) \frac{|\nabla P(x)|^2}{P^2(x)} dx \leq 4 \int_{\mathbb{R}^d} |\nabla u(x)|^2 dx.$$

Lemma 0.6 *The Sobolev space $\dot{H}^1(\mathbb{R}^d)$ be invariant under the transformation :*

$$u \rightarrow \tilde{u} = e^{iw}u, \quad v \rightarrow \tilde{v} = e^{iw}v,$$

$$\|u\|_{\dot{H}^1} \simeq \|\tilde{u}\|_{\dot{H}^1}, \quad \|v\|_{\dot{H}^1} \simeq \|\tilde{v}\|_{\dot{H}^1}$$

Proof. Clearly, We observe that $v \in \dot{H}^1(\mathbb{R}^d)$

$$\nabla(e^{iw}u) = (iu\nabla w + \nabla u)e^{iw}.$$

Consequently, for every $u \in \dot{H}^1(\mathbb{R}^d)$,

$$\|\tilde{u}\|_{\dot{H}^1} = \|e^{iw}u\|_{\dot{H}^1} \leq \|u\nabla w\|_{L^2} + \|\nabla u\|_{L^2}.$$

We set

$$w = \log P\mu, \quad v = P\mu.$$

Note that

$$\nabla w = \frac{\nabla P}{P}.$$

Hence, by lemma 0.5,

$$\|u\nabla w\|_{\dot{H}^1} = \left\| u \frac{\nabla P}{P} \right\|_{\dot{H}^1} \leq 2\|u\|_{\dot{H}^1}.$$

From this, it follows

$$\|\tilde{u}\|_{\dot{H}^1} \leq 3\|u\|_{\dot{H}^1}.$$

Using similar estimates for $\tilde{u} \rightarrow u = e^{-iw}\tilde{u}$, we deduce

$$\frac{1}{3}\|u\|_{\dot{H}^1} \leq \|\tilde{u}\|_{\dot{H}^1}.$$

This completes the proof of lemma.

We now establish the main result of this paper.

Theorem 0.7 *Let $\vec{F} \in \mathcal{D}'(\mathbb{R}^d)$, $d \geq 3$. Suppose that, for all $u, v \in \mathcal{D}(\mathbb{R}^d)$*

$$\left| \left\langle \vec{F}, v\nabla \bar{u} - \bar{u}\nabla v \right\rangle \right| \leq C \|u\|_{\dot{H}^1} \|v\|_{\dot{H}^1}. \quad (2)$$

Then there is a vector-field $\vec{G} = \nabla \left(\Delta^{-1} \operatorname{div} \vec{F} \right) \in L_{loc}^2(\mathbb{R}^d)$ such that

$$\int_{\mathbb{R}^d} \left| \vec{G}(x) \right|^2 |u(x)|^2 dx \leq C \|u\|_{\dot{H}^1}^2$$

for all $u \in \mathcal{D}(\mathbb{R}^d)$, where the constant C does not depend on $u \in \mathcal{D}(\mathbb{R}^d)$.

Proof. Suppose that

$$\left| \left\langle \vec{F}, v \nabla \bar{u} - \bar{u} \nabla v \right\rangle \right| \leq C \|u\|_{\dot{H}^1} \|v\|_{\dot{H}^1}.$$

holds. By the continuity the bilinear form on the left-hand side can be extended to all $u, v \in \dot{H}^1(\mathbb{R}^d)$. Let v be a nonnegative function such that $w = \log v$ and by the proposition 0.1

$$\nabla w = \frac{\nabla v}{v} \in L_{loc}^2(\mathbb{R}^d).$$

We need w to be chosen so that the Sobolev space $\dot{H}^1(\mathbb{R}^d)$ be invariant under the transformation :

$$\begin{aligned} u &\rightarrow \tilde{u} = e^{iw} u, & v &\rightarrow \tilde{v} = e^{iw} v, \\ \|u\|_{\dot{H}^1} &\simeq \|\tilde{u}\|_{\dot{H}^1}, & \|v\|_{\dot{H}^1} &\simeq \|\tilde{v}\|_{\dot{H}^1} \end{aligned}$$

Applying lemma 0.6 and (2), with $\tilde{u}$ and $\tilde{v}$ in place of u and v , we get

$$\begin{aligned} \left| \left\langle \vec{F}, \tilde{v} \nabla \tilde{u} - \tilde{u} \nabla \tilde{v} \right\rangle \right| &\leq C \|\tilde{u}\|_{\dot{H}^1} \|\tilde{v}\|_{\dot{H}^1} \\ &\leq 9C \|u\|_{\dot{H}^1} \|v\|_{\dot{H}^1}. \end{aligned}$$

Notice that

$$\tilde{v} \nabla \tilde{u} - \tilde{u} \nabla \tilde{v} = v \nabla \bar{u} - \bar{u} \nabla v - 2i\bar{u}v \nabla w.$$

Combining the preceding estimates, we obtain

$$\begin{aligned} 2 \left| \left\langle \vec{F}, \bar{u} v \nabla w \right\rangle \right| &\leq \left| \left\langle \vec{F}, v \nabla \bar{u} - \bar{u} \nabla v \right\rangle \right| + \left| \left\langle \vec{F}, \tilde{v} \nabla \tilde{u} - \tilde{u} \nabla \tilde{v} \right\rangle \right| \\ &\leq 10C \|u\|_{\dot{H}^1} \|v\|_{\dot{H}^1}. \end{aligned}$$

Observe that

$$v \nabla w = \nabla v.$$

Thus, we arrive at the inequality

$$\left| \left\langle \vec{F}, \bar{u} \nabla v \right\rangle \right| \leq C' \|u\|_{\dot{H}^1} \|v\|_{\dot{H}^1}.$$

From the preceding estimate and (2), we deduce

$$\begin{aligned} \left| \left\langle \vec{F}, \nabla (\bar{u} v) \right\rangle \right| &= \left| \left\langle \vec{F}, v \nabla \bar{u} + \bar{u} \nabla v \right\rangle \right| \leq \left| \left\langle \vec{F}, v \nabla \bar{u} \right\rangle \right| + \left| \left\langle \vec{F}, \bar{u} \nabla v \right\rangle \right| \\ &\leq C'' \|u\|_{\dot{H}^1} \|v\|_{\dot{H}^1}. \end{aligned}$$

This yields

$$\left| \left\langle \left(\operatorname{div} \vec{F} \right) u, v \right\rangle \right| \leq C'' \|u\|_{\dot{H}^1} \|v\|_{\dot{H}^1},$$

where $u \in \dot{H}^1(\mathbb{R}^d)$ and $v = P\mu$. By theorem 2.2 in [MV],

$$\operatorname{div} \vec{F} \in \mathcal{M}(\dot{H}^1 \rightarrow \dot{H}^{-1}).$$

Hence, there is a vector-field $\vec{G} \in L_{loc}^2(\mathbb{R}^d)$ such that

$$\vec{G} = \nabla \left(\Delta^{-1} \operatorname{div} \vec{F} \right) = \nabla f \in \mathcal{M}(\dot{H}^1 \rightarrow L^2).$$

Corollary 0.8 *Under the assumptions of theorem 0.7, it follows that if*

$$\left| \left\langle \vec{F}, v \nabla \bar{u} - \bar{u} \nabla v \right\rangle \right| \leq C \|u\|_{\dot{H}^1} \|v\|_{\dot{H}^1},$$

then

$$\Delta f \in \mathcal{M}(\dot{H}^1 \rightarrow \dot{H}^{-1}).$$

Theorem 0.9 *Let $f \in \mathcal{D}'(\mathbb{R}^d)$. Then the inequality*

$$|\langle [f, \Delta] u, v \rangle| \leq C \|u\|_{\dot{H}^1} \|v\|_{\dot{H}^1},$$

holds for all $u, v \in \mathcal{D}(\mathbb{R}^d)$, if and only if there is a vector-field $\vec{G} \in L_{loc}^2(\mathbb{R}^d)$ such that

$$\int_{\mathbb{R}^d} \left| \vec{G}(x) \right|^2 |u(x)|^2 dx \leq C \|u\|_{\dot{H}^1}^2,$$

where $\vec{G} = \nabla f$.

Proof. Suppose that $\vec{G} = \nabla \left(\Delta^{-1} \operatorname{div} \vec{F} \right)$ and

$$\int_{\mathbb{R}^d} \left| \vec{G}(x) \right|^2 |u(x)|^2 dx \leq C \|u\|_{\dot{H}^1}^2.$$

Then $\vec{G} \cdot \nabla$ is form bounded, since by Schwarz's inequality, we have

$$\left| \left\langle \vec{G} \cdot \nabla u, v \right\rangle \right| \leq \left\| u \vec{G} \right\|_{L^2} \|\nabla v\|_{L^2} \leq C \|u\|_{\dot{H}^1} \|v\|_{\dot{H}^1}.$$

Hence, this inequality obviously yields

$$\begin{aligned} \left| \left\langle \vec{F}, v \nabla \bar{u} - \bar{u} \nabla v \right\rangle \right| &\leq \left| \left\langle \vec{F}, v \nabla \bar{u} \right\rangle \right| + \left| \left\langle \vec{F}, \bar{u} \nabla v \right\rangle \right| \\ &\leq C \|u\|_{\dot{H}^1} \|v\|_{\dot{H}^1}. \end{aligned}$$

This completes the proof of theorem.

References

- [LL] Lieb, E.H., and Loss, M.: Analysis, second Edition, Amer.math.Soc.Providence, RI, 2001.
- [MV] Maz'ya, V., Verbitsky, I. The Schrödinger operator on the energy space: Boundness and compactness criteria, Acta Mathematica, 188 (2002), 263-302.